


INCLUSIVE DIRECT CP-ASYMMETRIES IN CHARMLESS $B^\pm$ -DECAYS^a

Ulrich Nierste

DESY - Theory group, Notkestrasse 85, D-22607 Hamburg, Germany



Direct CP-asymmetries in inclusive decay modes can be cleanly calculated by exploiting quark-hadron duality. This is in sharp contrast to CP-asymmetries in exclusive channels, where unknown strong phases prevent a clean extraction of CKM parameters from measured CP-asymmetries. We have calculated the inclusive CP-asymmetries in $B^\pm$ -decays into charmless final states with strangeness one or strangeness zero. In our results large logarithms are properly summed to all orders. We find

$$a_{CP}(\Delta S = 0) = \left(2.0^{+1.2}_{-1.0}\right) \%, \quad a_{CP}(\Delta S = 1) = (-1.0 \pm 0.5) \%.$$

The constraints on the apex $(\bar{\rho}, \bar{\eta})$ of the unitarity triangle obtained from these two CP-asymmetries define circles in the $(\bar{\rho}, \bar{\eta})$ -plane. $a_{CP}(\Delta S = 0)$ measures $\sin \gamma \cdot |V_{cb}/V_{ub}|$. The presented work has been done in collaboration with Gaby Ostermaier and Alexander Lenz.

First we define the average branching fraction in terms of the decay rate Γ :

$$\overline{Br} = \frac{\Gamma(B^+ \rightarrow X) + \Gamma(B^- \rightarrow \bar{X})}{2\Gamma_{tot}}. \quad (1)$$

In the following we are interested in inclusive final states X containing no charmed particle. Further the total strangeness S of X must be known, we will consider the cases $X = X(S = 0)$ and $X = X(|S| = 1)$. Similarly we define the direct CP-asymmetries as

$$A_{CP} = \frac{1}{2} \left[Br(B^+ \rightarrow X) - Br(B^- \rightarrow \bar{X}) \right], \quad a_{CP} = \frac{A_{CP}}{\overline{Br}}. \quad (2)$$

The measurement of $\overline{Br}$ in (1) and of the CP-asymmetries in (2) requires a sum over semi-inclusive final states in which the Kaons and strange baryons must be identified. Direct CP-asymmetries in exclusive decay modes are hard to access theoretically. Non-perturbative rescattering effects induce strong phases, which are difficult to estimate. On the contrary for the case of inclusive final states local quark-hadron duality allows to calculate the quantities in (1) and (2) reliably within perturbation theory. Theorists have been considering inclusive direct CP-asymmetries since 1979¹. Up to now the inclusive a_{CP} 's were

^aTalk given at the *Recontre de Moriond* on QCD and High Energy Hadronic Interactions, March 21st to 28th, 1998, Les Arcs, France, to appear in the proceedings.




Figure 1: The lightly (darkly) shaded area shows the constraint on $(\bar{\rho}, \bar{\eta})$ stemming from $a_{CP}(\Delta S = 0)$ ($a_{CP}(\Delta S = 1)$), if the a_{CP} 's are measured as in (3).

believed to be small, of the order of a few permille. Yet in the predictions of¹ either large logarithms have not been summed to all orders or m_t was taken too small. This has been corrected for in our new paper², in which also the predictions for charmless branching ratios calculated before³ have been improved by incorporating new QCD corrections. We find

$$a_{CP}(\Delta S = 0) = \left(2.0^{+1.2}_{-1.0}\right) \%, \quad a_{CP}(\Delta S = 1) = (-1.0 \pm 0.5) \%.$$

Here the error bars stem from the uncertainty in m_c/m_b and $\bar{\rho}, \bar{\eta}$ and from the residual dependence on the renormalization scale μ . The μ -dependence can be reduced by calculating certain two-loop diagrams. This is possible with reasonable effort and will be done, once the inclusive direct CP-asymmetries receive experimental interest.

The dependence of the a_{CP} 's on $\bar{\rho}$ and $\bar{\eta}$ is welcome in order to constrain the apex of the unitarity triangle. For a model scenario with

$$a_{CP}(\Delta S = 0) = 2.0 \%, \quad a_{CP}(\Delta S = 1) = -1.0 \% \quad (3)$$

and an assumed total error of 20 % the constraints on $\bar{\rho}, \bar{\eta}$ are shown in figure 1. They are nice circles in the $\bar{\rho}, \bar{\eta}$ -plane, whose information is complementary to the familiar circle from B - $\bar{B}$ -mixing and the hyperbola from ε_K^A . The A_{CP} 's defined in (2) are simply proportional to $\bar{\eta}$.

References

1. M. Bander, D. Silverman and A. Soni, Phys. Rev. Lett. 43 (1979) 242; J.-M. Gérard and W.-S. Hou, Phys. Rev. D43 (1991) 2909; H. Simma, G. Eilam and D. Wyler, Nucl. Phys. B352 (1991) 367; L. Wolfenstein, preprint no. NSF-ITP-90-29 (unpublished), and Phys. Rev. D43 (1991) 151; Yu. Dokshitser and N. Uraltsev, JETP. Lett. 52 (1990) 509; N. Uraltsev, hep-ph/9212233.
2. A. Lenz, U. Nierste and G. Ostermaier, DESY-97-208, hep-ph/9802202.
3. A. Lenz, U. Nierste and G. Ostermaier, Phys. Rev. D56 (1997) 7228.
4. A. J. Buras, M. Jamin and P. H. Weisz, Nucl. Phys. B347 (1990) 491; S. Herrlich and U. Nierste, Phys. Rev. D52 (1995) 6505.